

1 Mass Spectrometry in Bioprospecting of Plant Natural Products in the Amazon

2 Rainforest

3 *L. Soto*

4 Abstract

5 The Amazon rainforest is a megadiverse region of cultural
6 complexity. Ethnic groups have been using an extensive plant
7 repository for several applications. Bioprospecting the potential
8 bioactive molecules requires a set of analytical techniques capable of
9 elucidating the chemical composition of complex samples.
10 Chromatography coupled with mass spectrometry is one of the most
11 accurate and robust strategies available for identifying and quantifying natural products. This brief
12 review outlines the most common mass spectrometry techniques and applications in plant
13 bioprospecting in the Amazon region during the last ten years.



Source: ³

14 **Keywords:** bioprospecting, natural products, mass spectrometry, LC-MS, GC-MS, Amazon.

15 Content

16		
17	1. Introduction	2
18	2. Methodology.....	3
19	3. MS techniques	3
20	GC-MS.....	3
21	LC-MS Orbitrap.....	6
22	LC-MS TOF.....	7
23	MALDI TOF.....	7
24	4. Alternative approaches	7
25	In-vivo SPME and LC-MS QqQ	7
26	Off-line counter-current chromatography.....	8
27	5. Data processing.....	8
28	6. Conclusion and future perspectives.....	8
29	7. References	9

30

1. Introduction

The Amazon rainforest covers around 5 500 000 km² across nine South American countries, mainly Brazil. The tropical rainforest is rich in biome, constituting the world's largest collection of flora and fauna. This biome houses millions of insect species, thousands of plants, birds, and mammals. Some approximations indicate that the Amazon host around 16 000 tree species,¹ and that 0.25 km² of Ecuadorian rainforest contains more than 1 100 species of trees.²

The Amazon is also culturally complex. It includes thousands of indigenous territories and more than 30 million people and 320 different ethnic groups.³ These communities have their own traditional norms and knowledge that shapes the social organization and their interaction with the environment. Over generations, they transferred traditional medicine that employs natural resources to treat various health conditions.

The Amazon region has a proven application of natural products and ethnobotanical surveys play a considerable role in bioprospecting new bioactive compounds. An ethnobotanical study for collecting data on medicinal plants in the Valley of Jurueña, Brazil, found 332 plant species of 90 families reported by interviewed people. Those plants were used for various gastrointestinal and respiratory ailments⁴. In 2015, another ethno-directed study with the quilombola communities in Oriximiná, Brazil, examined the activity of 11 plants with *in vitro* cultures parasites for malaria. The results showed the antiplasmodial potential of *Aspidosperma rigidum* and *Bertholletia excelsa*⁵.

Chromatography, mass spectrometry (MS), and nuclear magnetic resonance (NMR), are the most common analytical tools for exploring natural products. Mass spectrometry is widely used in diverse fields to identify and quantify the chemical composition of complex samples. Mass spectrometry is a rapid, sensitive, and accurate method that measures the mass-to-charge ratio of analytes. Fragmentation of molecules with tandem mass spectrometry can provide more relevant information on the chemical structure. Along with mass spectrometry, chromatographic separation can enhance selectivity. The retention time of the compounds gives an additional identity indication.

This review aims to explore the current mass spectrometry techniques applied to the bioprospection of natural products from plants in the Amazon region.

2. Methodology

In this non-systematic review, most of the articles were obtained from the ScienceDirect database with the query “((bioprospecting OR bioprospection) AND (amazon) AND (mass AND spectrometry OR ms OR hrms))”. The publication period was 2012-2022. Additionally, searches from Scopus, Google Scholar, Web of Science, and specialized journals were included. The articles are mainly in English. Other languages, such as Portuguese and Spanish, could be considered to explore other databases in a further systematic study.

3. MS techniques

The most common analytical technique was gas chromatography (GC) coupled to single quadrupole mass analyzer, followed by other liquid chromatography (LC) coupled to Orbitrap, time-of-flight, and triple quadrupole analyzers. Nuclear magnetic resonance (NMR) was also used in some studies for identity confirmation.

The predominance of GC-MS single quadrupole over other techniques could be explained by the scope of applications and its popularity. The scope of many studies was on investigating small and volatile molecules, while the heavier and thermo-labile ones were not considering for which other techniques are usually necessary. The GC-MS single quadrupole instrument is a standard instrument in many analytical laboratories and used in routine analysis.

Besides, it is worth mentioning that the findings are limited to the articles in this review and the way in how the articles were collected, mainly via the search query described in the methodology.

GC-MS

Gas chromatography-mass spectrometry (GC-MS) has been extensively used in plant bioprospecting of essential oils and volatile compounds. GC-MS was the most common technique found in this review, and it could be due to the specific application scope to volatile analytes and that it is commonly available. For the analysis, samples are usually taken from aerial parts, for instance, leaves and small stems. Some of the analytes were small molecules such as essential oils, terpenoids, and alkaloids. For certain plant species, these compounds and the chemical composition were reported for the first time.


87 GC-MS is a well-studied and standardized technique. In all GC-MS studies, the electron impact
88 ionization was 70 eV. The most common mass analyzer was the single quadrupole, and the mass
89 range was approximately between 35 and 500 m/z. The mass spectra are compared with spectra
90 libraries and specialized books. For qualitative analysis and relative quantification, most studies
91 used the linear retention index with Van Den Dool and Kratz equation, series of n-alkenes C8-
92 C38, along with NIST and FFNSC libraries ^{6 7 8 9}. While for quantitative analysis, one study
93 prepared an external calibration. ¹⁰ Herein, some details about the applications of this technique
94 are presented.

95 GC-MS single quadrupole has been of particular interest to study essential oils. Samples from
96 *Diospyros guianensis*, *Carapa guianensis* and *Aspidosperma nitidum* showed larvicidal activity
97 ¹¹. In order to determine the bioactive compounds, GC-MS analysis has been helpful in the
98 identification of some constituents, for example, fatty acids, limonoids, and plumbagin. A possible
99 synergic effect of some of these compounds was also suggested. For some compounds, the
100 structure elucidation additionally involved NMR and LC-MS.

101 The seasonal variation of the chemical composition of some essential oils has been studied. The
102 *Mesosphaerum suaveolens* (L.) Kuntze, an endemic plant from the Brazilian Amazon, has a
103 different composition of essential oils during the rainy, intermediate, and dry seasons, as shown in
104 *Figure 1*. ⁶ The dry season showed a greater linalool presence than other seasons. Contrary to this
105 finding, Pimentel et al. ¹⁰ found a greater concentration of linalool in the wet season for another
106 plant, *Aniba rosaeodora* Ducke. This suggests that essential oils are not directly influenced by
107 environmental factors but intrinsic ones. Pimentel et al. ¹⁰ also found greater effectiveness against
108 phytopathogens for those essential oils that were extracted during the wet season. The study also
109 suggested linalool as the main active compound. In a similar study, the larvicidal activity of
110 essential oils from *E. uniflora*, *L. camara*, *O. basilicum*, *P. neochilus* against *A. aegypti* showed
111 greater effectiveness in the dry season ⁷.

112 Geographical and environmental factors could also influence the chemical profile of the plants.
113 Plant samples that were collected at different locations differed in their essential oils composition
114 ⁸. Seasons can also modify the overall extraction yield and composition. Lower volatile
115 compounds can be explained by a higher sun radiation exposure⁶. An examination of chemical

116 profile against these and other factors could help to determine the best conditions for material
 117 collection depending on the compounds of interest.



118
 119 **Figure 1** Essential oil composition variation over the seasons for *Mesosphaerum suaveolens*.
 120 Figure from Luz et al. ⁶

121 The shell of *Carapa guianensis* Aubl., a plant that could be a potential eco-friendly larvicide,
 122 showed a possible synergic effect of fatty acids and limonoids. A mixture of essential oil extracts
 123 has been used to study the combined effect on bioactivity ¹¹.

124 Potential natural products against helminth infections have also been reported in the context of
 125 growing resistance to common drugs. For example, a study of anthelmintic activity of the leaves
 126 and fine stems of *Hyptis dilatata* and *Mesosphaerum suaveolens* showed that the oils, containing
 127 terpenes, can have a synergistic action as nematocidal ⁸. The effect of monoterpenes has been
 128 attributed to their interference with receptors or enzymes ⁸. However, the exact mechanism of
 129 terpenes in biological systems is not yet well understood.

130 Some mosses have been traditionally used as medicinal plants ¹². GC-MS has helped to examine
 131 their chemical composition. For the study of *Neckeropsis undulata* (Hedw.) Reichardt, distillation-
 132 extraction, GC, and GC-MS single quadrupole were applied⁹. Some compounds were 1-octen-3-
 133 ol, α-muuirolol, naphthalene, and n-hexanal. The relative quantitation of each compound was
 134 calculated by peak normalization in GC coupled to a flame ionization detector (FID) ⁹. In a similar
 135 study for *Calymperes palisotii* Schwaegr, 13 compounds were identified and relatively quantified
 136 by GC/MS single quadrupole. 3-methyl-2-pentanone, phenylacetaldehyde, and E-nerolidol were
 137 the major constituents. E-nerolidol and furfural were first reported for mosses¹³. Epizonarene, α-
 138 selinene, β-selinene have been also identified as constituents of mosses oil extracts ¹⁴.

The smokes of oleoresin have been attributed positive effects for headaches. The smoke from *Burseraceae pitch (breu)* consisted mainly of terpenoids that were transferred to the smoke by volatilization. GC-MS single quadrupole identified at least 32 compounds¹⁵.

New compounds are usually reported for the first time in some studies, for example, new compounds in the family *Verbenaceae*¹⁶, and secondary metabolites for *E. amazonica* and *M. dubia*¹⁷. At the same time, some authors suggest that the presence of specific molecules could be attributed to other organisms¹³.

LC-MS Orbitrap

Unlike GC-MS, mainly limited to studying small, thermostable, and volatile compounds, liquid chromatography-mass spectrometry (LC-MS) opened a wide range of discovery possibilities. Non-volatile chemical compositions have not been explored yet for some medicinal plants as these molecules may require specific sample preparations, derivatization or cannot be determined by GC-MS. For this analysis, some of the samples were leaves, stems, bark, and seeds. The most common ion source interphase between LC and the mass spectrometer was electrospray ionization (ESI).

Some studies used the Orbitrap mass analyzer. An extensive analysis of the potential antimicrobial activity of 59 plant species from 33 families was carried out. 12 plants showed antibacterial activity. The study also isolated and identified three secondary metabolites for the first time. The analytical methods were chromatography, NMR, and high-resolution mass spectrometry (HRMS). An MS analysis was done with ESI in positive and negative mode with Orbitrap in the mass range of 100 to 850 Da. Alkaloid and triterpenic compounds were identified¹⁷. In another study, a hybrid mass analyzer linear trap quadrupole (LTQ) - Orbitrap was used to identify flavonoids in *Lippia origanoides* (Verbenaceae), a plant known for its medicinal and food applications. The ESI source was in positive and negative mode. In this case, the resolution was 30000, and the mass ranged from 140 to 1500 m/z¹⁶. For fragmentation, a data dependent scan was set with a collision energy of 35 eV and precursor width of 2.0 m/z¹⁶.

Ampelozizyphus amazonicus Ducke is a plant with medicinal attributes used in the Amazon region. In a study, an extract was analyzed by HPLC-HRMS. In this case, the ESI source was in negative mode. The Orbitrap allowed a resolution of 60000 for the full mass and 7500 for the data-

dependent acquisition. The collision energy for the collision-induced dissociation (CID) was 35 eV, and the width for the precursor ion was 2.0 m/z. For identification, the fragmentation patterns were relevant to propose several saponin chemical structures ¹⁸.

LC-MS TOF

A few studies included time-of-flight (TOF) analyzers. Spilanthol content and bactericide/antibiofilm activity have been determined for the *Acmella oleracea*, a plant with anti-cariogenic properties. HPLC-MS/MS with a hybrid Q-TOF (quadrupole-TOF) mass analyzer was used to quantify the concentration of spilanthol in the leaves. The ESI operated in positive mode and the range was from 50 to 1000 m/z ¹⁹. In another study of *Carapa guianensis*, a plant with potential eco-friendly larvicidal activity, HPLC-MS was used with ESI and Q-TOF to analyze the seed shells ¹¹.

MALDI TOF

The TOF analyzer is well suited to the pulse nature of laser desorption of matrix-assisted laser desorption/ionization (MALDI) and could represent an opportunity to explore large, unstable, and polymeric biomolecules by direct infusion. Research papers using MALDI-TOF are even less common. MALDI – TOF in positive mode has been used to determine the molecular weight of a phenolic compound (761.2 Da) with antioxidant activity from epicarp samples of *Garcinia brasiliensis*, a plant with attributed anti-inflammatory, antioxidant and antitumor properties ²⁰.

4. Alternative approaches

In-vivo SPME and LC-MS QqQ

An in-vivo solid phase microextraction (SPME) sampling technique has been documented. Twenty Amazonian plant species were sampled in vivo using SPME probes. Probes are inserted under the plant's bark, left for a contact time, then transferred to plastic tubes under vacuum. Samples are extracted from the probes with ethanol solution, evaporated, and reconstituted before the analysis. With this approach, the damage to the plant is minimized, the sampling is faster than the standard preparation procedures, and it could be suitable for sampling covering large remote areas. For the LC-MS analysis, this study used ESI along a triple quadrupole (QqQ) mass analyzer for the range 150 - 600 m/z. ²¹




Figure 2 Graphical abstract for in-vivo sampling with SPME probe. Image from Musteata et al.²¹

Off-line counter-current chromatography

Counter-current chromatography (CCC) has been reported to improve the identification of constituents in complex samples as a purification step prior to mass spectrometry analysis. In a study of the phenolic profile of *Lippia origanoides*, counter-current chromatography was used as a preparatory step for fractionation of a sample extract. This approach increased the number of identified compounds¹⁶. CCC has also been used to fractionate *Garcinia brasiliensis* epicarp extract samples²⁰.

5. Data processing

MS spectra is compared to chemo-taxonomic data available in literature and databases. Some studies used the software Xcalibur, and a common mass spectra library was NIST. Open-source processing software, e.g. MZMine, or collaborative mass spectra annotation, e.g. Global Natural Products Social Molecular Networking (GNPS), seem to be not common in these studies. Multivariate analysis and a machine learning algorithm has been applied. Partial least squares - discriminant analysis (PLS-DA) and random forest analysis helped to differentiate between two types of oleoresins and find a potential chemical marker based on mass spectrometry data²².

6. Conclusion and future perspectives

Mass spectrometry experiments offer a variety of possibilities for bioprospecting. This brief review outlined some of the most common mass spectrometry techniques and their applications in bioprospecting plants in the Amazon region. It includes an overview of GC-MS, LC-MS, Orbitrap, TOF, and MALDI, and describes the typical samples, analytes, technique settings, and data processing. This review is limited to the revised articles here and the way in which the articles were collected, mainly via the search query. GC-MS single quadrupole was found to be the most common technique. This could be explained because of the intended application to identify or

quantify small molecules e.g., essential oils, and the greater popularity of this instrument. The main challenge remains in expanding the analytical capabilities offered by MS to analyze a broader spectrum of natural products. In this regard, high-resolution mass spectrometry, ambient mass spectrometry, mass spectrometry imaging and other ionization modes could lead to a better understanding of the complex chemical configuration of plants in the Amazon region and their potential applications.

7. References

1. ter Steege H, Pitman NCA, Sabatier D, et al. Hyperdominance in the Amazonian Tree Flora. *Science*. 2013;342(6156):1243092. doi:10.1126/science.1243092
2. Wright JS. Plant diversity in tropical forests: a review of mechanisms of species coexistence. *Oecologia*. 2002;130(1):1-14. doi:10.1007/s004420100809
3. Thompson C. *Amazon Alive! A Decade of Discovery 1999-2009*. WWF Living Amazon Initiative https://wwfint.awsassets.panda.org/downloads/amazon_alive_web_ready_sept23.pdf
4. Bieski IGC, Leonti M, Arnason JT, et al. Ethnobotanical study of medicinal plants by population of Valley of Juruena Region, Legal Amazon, Mato Grosso, Brazil. *J Ethnopharmacol*. 2015;173:383-423. doi:10.1016/j.jep.2015.07.025
5. Oliveira DR, Krettli AU, Aguiar ACC, et al. Ethnopharmacological evaluation of medicinal plants used against malaria by quilombola communities from Oriximiná, Brazil. *J Ethnopharmacol*. 2015;173:424-434. doi:10.1016/j.jep.2015.07.035
6. Luz TRSA, Leite JAC, de Mesquita LSS, et al. Seasonal variation in the chemical composition and biological activity of the essential oil of *Mesosphaerum suaveolens* (L.) Kuntze. *Ind Crops Prod*. 2020;153:112600. doi:10.1016/j.indcrop.2020.112600
7. Luz TRSA, Leite JAC, de Mesquita LSS, et al. Seasonal variation in the chemical composition and larvicidal activity against *Aedes aegypti* L. of essential oils from Brazilian Amazon. *Exp Parasitol*. 2022;243:108405. doi:10.1016/j.exppara.2022.108405
8. Lima AS, Fernandes YML, Silva CR, et al. Anthelmintic evaluation and essential oils composition of *Hyptis dilatata* Benth. and *Mesosphaerum suaveolens* Kuntze from the Brazilian Amazon. *Acta Trop*. 2022;228:106321. doi:10.1016/j.actatropica.2022.106321
9. Miranda TG, Alves RJM, de Souza RF, Maia JGS, Figueiredo PLB, Tavares-Martins ACC. Volatile concentrate from the neotropical moss *Neckeropsis undulata* (Hedw.) Reichenhardt, existing in the Brazilian Amazon. *BMC Chem*. 2021;15(1):7. doi:10.1186/s13065-021-00736-3
10. Pimentel RBQ, Souza DP, Albuquerque PM, et al. Variability and antifungal activity of volatile compounds from *Aniba rosaeodora* Ducke, harvested from Central Amazonia in two different seasons. *Ind Crops Prod*. 2018;123:1-9. doi:10.1016/j.indcrop.2018.06.055
11. Correa de Oliveira PM, Barreto Sousa JP, Albernaz LC, Coelho-Ferreira M, Salmen Espindola L. Bioprospection for new larvicides against *Aedes aegypti* based on ethnoknowledge from the

- 257 Amazonian São Sebastião de Marinaú riverside community. *J Ethnopharmacol.* 2022;293:115284.
258 doi:10.1016/j.jep.2022.115284
- 259 12. Asakawa Y, Ludwiczuk A, Nagashima F. Phytochemical and biological studies of bryophytes.
260 *Phytochemistry.* 2013;91:52-80. doi:10.1016/j.phytochem.2012.04.012
- 261 13. Alves RJM, Miranda TG, Pinheiro RO, Andrade EHDA, Tavares-Martins ACC. Analysis of volatile
262 organic compounds from the moss *Calymperes palisotii* Schwaegr. *South Afr J Bot.* 2022;150:974-
263 977. doi:10.1016/j.sajb.2022.09.006
- 264 14. Valarezo E, Vidal V, Calva J, Jaramillo SP, Febres JD, Benitez A. Essential Oil Constituents of Mosses
265 Species from Ecuador. *J Essent Oil Bear Plants.* 2018;21(1):189-197.
266 doi:10.1080/0972060X.2018.1432420
- 267 15. Albino R da C, Braz MM, Bizzo HR, Santana da Silva RV, Leitão SG, Ribeiro de Oliveira D.
268 Amazonian medicinal smokes: Chemical analysis of Burseraceae pitch (breu) oleoresin smokes and
269 insights into their use on headache. *J Ethnopharmacol.* 2021;276:114165.
270 doi:10.1016/j.jep.2021.114165
- 271 16. Leitão SG, Leitão GG, Vicco DKT, et al. Counter-current chromatography with off-line detection by
272 ultra high performance liquid chromatography/high resolution mass spectrometry in the study of the
273 phenolic profile of *Lippia origanoides*. *J Chromatogr A.* 2017;1520:83-90.
274 doi:10.1016/j.chroma.2017.09.004
- 275 17. Roumy V, Ruiz Macedo JC, Bonneau N, et al. Plant therapy in the Peruvian Amazon (Loreto) in case
276 of infectious diseases and its antimicrobial evaluation. *J Ethnopharmacol.* 2020;249:112411.
277 doi:10.1016/j.jep.2019.112411
- 278 18. Simen TJM, Finotelli PV, Barboza FF, et al. Spray-dried extract from the Amazonian adaptogenic plant
279 *Ampelozizyphus amazonicus* Ducke (Saracura-mirá): Chemical composition and immunomodulatory
280 properties. *Food Res Int.* 2016;90:100-110. doi:10.1016/j.foodres.2016.10.040
- 281 19. Peretti P, Rodrigues ET, de Souza Junior BM, et al. Spilanthol content of *Acmella oleracea* subtypes
282 and their bactericide and antibiofilm activities against *Streptococcus mutans*. *South Afr J Bot.*
283 2021;143:17-24. doi:10.1016/j.sajb.2021.08.001
- 284 20. Gontijo VS, de Souza TC, Rosa IA, et al. Isolation and evaluation of the antioxidant activity of phenolic
285 constituents of the *Garcinia brasiliensis* epicarp. *Food Chem.* 2012;132(3):1230-1235.
286 doi:10.1016/j.foodchem.2011.10.110
- 287 21. Musteata FM, Sandoval M, Ruiz-Macedo JC, Harrison K, McKenna D, Millington W. Evaluation of
288 in vivo solid phase microextraction for minimally invasive analysis of nonvolatile phytochemicals in
289 Amazonian plants. *Anal Chim Acta.* 2016;933:124-133. doi:10.1016/j.aca.2016.05.053
- 290 22. Albino R da C, Simas RC, da Silva Martins K, et al. Differentiation of black and white pitch
291 (Burseraceae) oleoresins: A mass spectrometry-based chemoethnotaxonomic study. *J*
292 *Ethnopharmacol.* 2020;259:112968. doi:10.1016/j.jep.2020.112968